

Ollscoil na hÉireann, Gaillimh GX 2159
National University of Ireland, Galway
Summer Examinations, 2003/2004

Exam Code(s)	<u>1BA1;1BS1;1EL1;1ER1;1MR1;1PT1.</u>
Exam(s)	<u>First Year and First Arts</u>
Module Code(s)	<u>AM100</u>
Module(s)	<u>Applied Mathematics</u>
Paper No	<u>2</u>
Repeat Paper	<u>Special Paper</u>
External Examiner(s)	<u>Professor Brian Straughan</u>
Internal Examiner(s)	<u>Dr. M. Ó Confhaola</u>
	<u>Dr. P.M. O'Leary.</u>

Instructions:

Attempt **SIX** questions. You may assume throughout that the Earth's gravitational acceleration is $g = 10ms^{-2}$.

Duration	<u>3 hrs</u>
No. of Answer books	<u> </u>
Requirements	<u> </u>
Handout	<u> </u>
MCQ	<u> </u>
Statistical Tables	<u>Yes - Log Tables</u>
Graph paper	<u> </u>
Log Graph Paper	<u> </u>
Other Material	<u> </u>
No. of Pages	<u>3</u>
Department(s)	<u>Mathematical Physics</u>

1. A lift descends from rest with acceleration $k \text{ ms}^{-2}$, and after reaching a speed of 10 m/s continues at constant speed for 10s . It is brought to rest with a deceleration of $2k \text{ ms}^{-2}$. If the total time taken descending was 25s ,
 - a. Find the value of k ;
 - b. Find the distance travelled during the descent.
2. A block of mass 3kg rests on a horizontal table. It is attached by means of a light, inextensible string to a particle of mass 8 kg . The string passes over a smooth pulley at the edge of the table. There is a frictional force of 15N opposing the motion of the block. Find
 - a. the acceleration of the system;
 - b. the tension in the string.
3. A ball is kicked with a velocity of 15 m/s at an angle of 30° to the horizontal towards a wall which is 8 m away.
 - a. How far up the wall does the ball hit ?
 - b. What is the speed of the ball when it hits the wall ?
4. A ship A is moving due east at 18 km/h and another ship B is moving in a direction $\text{N } 120^\circ \text{ east}$ at 10 km/h . B is 12 km due north of A and both ships continue with the same velocities. Find the time when the ships are closest to each other and the shortest distance between them.
5. A 100 litre tank contains 20% pollutant. At time $t = 0$, a solution containing 10% pollutant flows into the tank at a rate of 5 litres/minute . The well-stirred mixture is drained off at the same rate.
 - a. Find how much pollutant there is in the tank after 10 minutes ;
 - b. How long until there is 15 litres of pollutant in the tank ?
 - c. How much pollutant will be in the tank as $t \rightarrow \infty$?
6. Find the position of the centre of mass of a uniform solid hemisphere of radius a .
 Prove that the centre of mass of a uniform hemispherical shell, whose inner and outer radii are a and b , is at a distance

$$\frac{3(a+b)(a^2+b^2)}{8(a^2+ab+b^2)}$$

from the centre and deduce the position of the centre of mass of a thin hemispherical shell.

7. A car is travelling round a circular bend of radius 25 m. The coefficient of friction between the car and the road is 0.3. The car has a mass of 500 kg. What is the maximum safe speed for the car if
- the road is unbanked
 - the road is banked at 15° to the horizontal ?
8. Tidal motion may be considered to be simple harmonic motion with a period of 12 hours. A certain tide has a low water mark of 5 m and a high water mark of 9 metres. Low tide occurs at 11.30 a.m.
- At what time may a ship requiring a draught of 6 m of water enter the harbour safely?
9. Two identical small spheres of mass 2 kg collide. The first, A , moves with a speed of 6 m/s and the second B with a speed of 8 m/s. Their directions of motion make angles of 30° and 45° respectively with their line of centres at the instant of contact. If the coefficient of restitution between the spheres is 0.4 find
- the velocities of the spheres after collision;
 - the kinetic energy lost during the collision.
10. A uniform ladder, of weight W , rests with one end against a smooth vertical wall and the other on a smooth horizontal floor. Its lower end is attached by means of a light inextensible string to the junction of the wall and the floor. A man of weight W climbs the ladder. Show that as the man moves from a point one-sixth of the way up the ladder to a point five-sixths of the way up, the tension in the string doubles.